

LE MATEMATICHE

Vol. LXVIII (2013) – Fasc. I, pp. 83–86

doi: 10.4418/2013.68.1.7

## A COMMENT ON COPOSINORMAL OPERATORS

H. CRAWFORD RHALY JR.

*Dedicated to Eric Xavier Mader*

Hyponormal operators are necessarily posinormal, but they need not be coposinormal. Coposinormality can nevertheless sometimes be an aid in determining hyponormality. This idea is applied to coposinormal factorable matrices whose interrupter is diagonal. Examples are given, including some Toeplitz matrices and some terraced matrices associated with the logistic sequence.

### 1. Introduction

If  $B(H)$  denotes the set of all bounded linear operators on a Hilbert space  $H$ , then  $M \in B(H)$  is *posinormal* if there exists a positive operator  $P \in B(H)$  satisfying  $MM^* = M^*PM$ . The operator  $P$  is referred to as the *interrupter*. From [2], Theorem 2.1, we know that  $M$  is posinormal if and only if

$$\lambda^2 M^* M \geq MM^*$$

for some  $\lambda \geq 0$ .  $M$  is *hyponormal* when  $\lambda = 1$ .  $M$  is *coposinormal* if  $M^*$  is posinormal. More information on posinormal operators can be found in [1], [2].

Some previous efforts to identify hyponormal operators had been focused on using posinormality in hopes that the interrupter  $P$  could be shown to satisfy

---

Entrato in redazione: 19 maggio 2012

AMS 2010 Subject Classification: 47B99.

Keywords: Posinormal operator, Hyponormal operator, Factorable matrix, Terraced matrix, Toeplitz matrix, Logistic sequence.

$I \geq P$ ; see [2], [3], [4]. This approach seemed reasonable and natural since all hyponormal operators are posinormal. Although hyponormal operators are not necessarily coposinormal, the following theorem tells us that there will be situations where coposinormality can also help us identify hyponormal operators.

**Theorem 1.1.** *Suppose that  $M \in B(H)$  is coposinormal; that is,  $M$  satisfies  $M^*M = MQM^*$  for some positive operator  $Q \in B(H)$ . If  $P$  is a positive operator and  $Q \geq P \geq I$ , then  $Y := \sqrt{P}M\sqrt{P}$  is hyponormal.*

*Proof.* If  $[Y^*, Y] := Y^*Y - YY^*$ , then

$$\langle [Y^*, Y]f, f \rangle = \langle (P - I)M\sqrt{P}f, M\sqrt{P}f \rangle + \langle (Q - P)M^*\sqrt{P}f, M^*\sqrt{P}f \rangle \geq 0$$

for all  $f$  in  $\ell^2$ , so  $Y$  is hyponormal.  $\square$

**Corollary 1.2.** *If  $M \in B(H)$  satisfies  $M^*M = MQM^*$  for some positive operator  $Q \in B(H)$  such that  $Q \geq I$ , then  $M$  is hyponormal.*

In the next section we apply this theorem and its corollary to lower triangular factorable matrices  $M$  acting on  $\ell^2$ .

## 2. Coposinormal Factorable Matrices with a Diagonal Interrupter

A lower triangular infinite matrix  $M = M(\{a_i\}, \{c_j\})$ , acting through multiplication to give a bounded linear operator on  $\ell^2$ , is *factorable* if its nonzero entries  $m_{ij}$  satisfy  $m_{ij} = a_i c_j$  where  $a_i$  depends only on  $i$  and  $c_j$  depends only on  $j$ ; if  $c_j = 1$  for all  $j$ , then  $M$  is *terraced*. Throughout this section we assume that  $M \in B(\ell^2)$  is a lower triangular factorable matrix.

**Theorem 2.1.** *Suppose  $M$  is a lower triangular factorable matrix that acts as a bounded operator on  $\ell^2$  and the following conditions are satisfied:*

- (1) both  $\{a_n\}$  and  $\{\frac{a_n}{c_n}\}$  are positive decreasing sequences that converge to 0;
- (2) the matrix  $T = [t_{mn}]$  by

$$t_{mn} = \begin{cases} \frac{a_m}{a_n} & \text{if } n = 0; \\ a_m \left( \frac{1}{a_n} - \frac{c_{n-1}}{c_n} \frac{1}{a_{n-1}} \right) & \text{if } 0 < n \leq m; \\ -\frac{c_{n-1}}{c_n} & \text{if } n = m + 1; \\ 0 & \text{if } n > m + 1. \end{cases}$$

is a bounded operator on  $\ell^2$ ;

- (3) the sequence  $\left\{ -\frac{c_{n-1}a_{n-1}}{c_n} + \left[ \frac{1}{a_n} - \frac{c_{n-1}}{c_n} \frac{1}{a_{n-1}} \right] \sum_{k=n}^{\infty} a_k^2 \right\}$  is constantly 0; and
- (4) the sequence  $\{c_n a_n\}$  is nonincreasing.

Then  $M$  is coposinormal with a diagonal interrupter, and furthermore,  $M$  is hyponormal.

*Proof.* As in [4, Theorems 3, 4], we have  $M = TM^*$ , so we compute the inter-rupter  $Q = T^*T$ ; the entries of  $Q = [q_{mn}]$  are given by

$$q_{mn} = \begin{cases} \frac{1}{a_0^2} \sum_{k=0}^{\infty} a_k^2 & \text{if } m = n = 0; \\ \frac{c_{n-1}^2}{c_n^2} + \left(\frac{1}{a_n} - \frac{c_{n-1}}{c_n} \frac{1}{a_{n-1}}\right)^2 \sum_{k=n}^{\infty} a_k^2 & \text{if } m = n \geq 1; \\ \frac{1}{a_0} \left(-\frac{c_{m-1}a_{m-1}}{c_m} + \left[\frac{1}{a_m} - \frac{c_{m-1}}{c_m} \frac{1}{a_{m-1}}\right] \sum_{k=m}^{\infty} a_k^2\right) & \text{if } m > n = 0; \\ \left(\frac{1}{a_n} - \frac{c_{n-1}}{c_n} \frac{1}{a_{n-1}}\right) \left(-\frac{c_{m-1}a_{m-1}}{c_m} + \left[\frac{1}{a_m} - \frac{c_{m-1}}{c_m} \frac{1}{a_{m-1}}\right] \sum_{k=m}^{\infty} a_k^2\right) & \text{if } m > n > 0; \\ \frac{1}{a_0} \left(-\frac{c_{n-1}a_{n-1}}{c_n} + \left[\frac{1}{a_n} - \frac{c_{n-1}}{c_n} \frac{1}{a_{n-1}}\right] \sum_{k=n}^{\infty} a_k^2\right) & \text{if } n > m = 0; \\ \left(\frac{1}{a_m} - \frac{c_{m-1}}{c_m} \frac{1}{a_{m-1}}\right) \left(-\frac{c_{n-1}a_{n-1}}{c_n} + \left[\frac{1}{a_n} - \frac{c_{n-1}}{c_n} \frac{1}{a_{n-1}}\right] \sum_{k=n}^{\infty} a_k^2\right) & \text{if } n > m > 0. \end{cases}$$

Inspection of the entries reveals that, because condition (3) is satisfied,  $Q$  will be a diagonal matrix;  $Q$  has the representation

$$Q = \text{diag} \left\{ \frac{\sum_{k=0}^{\infty} a_k^2}{a_0^2}, \frac{c_0 a_0}{c_1 a_1}, \frac{c_1 a_1}{c_2 a_2}, \dots, \frac{c_{n-1} a_{n-1}}{c_n a_n}, \dots \right\}.$$

Condition (4) then guarantees that  $Q \geq I$ , and it follows from Corollary 1.2 that  $M$  is hyponormal.  $\square$

**Example 2.2.** (Toeplitz Matrix) (a) Suppose  $M$  is the factorable matrix with entries  $m_{ij} = a_i c_j$  where  $a_i = r^i$ ,  $c_j = \frac{1}{r^j}$  for all  $i, j$  where  $0 < r < 1$ . If

$$Q := \text{diag} \left\{ \frac{1}{1-r^2}, 1, 1, 1, \dots \right\},$$

then  $MQM^* = M^*M$  is satisfied with  $Q \geq I$ , so  $M$  is hyponormal by Theorem 2.1.

(b) If  $D := \text{diag}\{d_n : n \geq 0\}$  where  $\frac{1}{1-r^2} \geq d_0 \geq 1$  and  $d_n = 1$  for  $n \geq 1$ , then  $\sqrt{D}M\sqrt{D}$  is also a hyponormal factorable matrix by Theorem 1.1.

**Example 2.3.** (a) For  $k > 0$ , let  $M$  denote the factorable matrix with entries  $m_{ij} = a_i c_j$  where  $a_i = \frac{1}{\sqrt{(i+k)(i+k+1)}}$  and  $c_j = \frac{\sqrt{j+k}}{\sqrt{j+k+1}}$  for all  $i, j$ . If

$$Q := \text{diag} \left\{ k+1, \frac{k+2}{k+1}, \frac{k+3}{k+2}, \dots, \frac{k+n+2}{k+n+1}, \dots \right\},$$

then  $MQM^* = M^*M$  is satisfied with  $Q \geq I$ , so  $M$  is hyponormal.

(b) If  $D := \text{diag}\{d_n : n \geq 0\}$  where  $k+1 \geq d_0 \geq 1$  and  $\frac{k+n+1}{k+n} \geq d_n \geq 1$  for  $n \geq 1$ , then  $\sqrt{D}M\sqrt{D}$  is also a hyponormal factorable matrix.

**Example 2.4.** (Recursion) (a) We consider the terraced matrix  $M$  determined as follows: Choose  $a_0, a_1$  such that  $0 < a_0 < 1$  and  $0 < a_1 \leq a_0(1-a_0)$ ; then

define  $b := \frac{(a_0 - a_1)}{a_0^2}$  and  $a_{i+1} = a_i(1 - ba_i)$  for each  $i \geq 1$ . We note that  $a_i \leq \frac{1}{i+1}$  for each  $i$ . Since  $\{a_i\}$  is decreasing, we have  $Q := \text{diag}\{\frac{1}{ba_0}, \frac{a_0}{a_1}, \frac{a_1}{a_2}, \dots\} \geq I$ ; finally, since  $MQM^* = M^*M$  is satisfied, it follows that  $M$  is a hyponormal operator on  $\ell^2$ . We note that in the special case when  $a_1 := a_0(1 - a_0)$ , we have  $b = 1$ , so we obtain the *logistic sequence*  $a_{i+1} = a_i(1 - a_i)$ , all  $i$ .

(b) If  $D := \text{diag}\{d_n : n \geq 0\}$  where  $\frac{1}{ba_0} \geq d_0 \geq 1$  and  $\frac{a_{n-1}}{a_n} \geq d_n \geq 1$  for  $n \geq 1$ , then  $\sqrt{DM}\sqrt{D}$  is another hyponormal factorable matrix.

## REFERENCES

- [1] C. S. Kubrusly - B. P. Duggal, *On posinormal operators*, Adv. Math. Sci. Appl. 17 (1) (2007), 131–147.
- [2] H. C. Rhaly Jr., *Posinormal operators*, J. Math. Soc. Japan 46 (4) (1994), 587–605.
- [3] H. C. Rhaly Jr., *Posinormal terraced matrices*, Bull. Korean Math. Soc. 46 (1) (2009), 117–123.
- [4] H. C. Rhaly Jr., *Posinormal factorable matrices whose interrupter is diagonal*, Mathematica (Cluj) 53 (76) (2) (2011), 181–188.

H. CRAWFORD RHALY JR.

1081 Buckley Drive,

Jackson, MS 39206, U.S.A.

e-mail: rhaly@member.ams.org